

PROPARGYL GLYCINE AMINO PROPARGYL DIOL COMPOUNDS  
FOR TREATMENT OF HYPERTENSION

RELATED APPLICATION

This application is a continuation-in-part of U.S. Application Serial No. 07/784,272 filed 29 October 1991.

FIELD OF THE INVENTION

Renin-inhibiting compounds are known for control of hypertension. Of particular interest herein are compounds useful as renin inhibiting agents.

BACKGROUND OF THE INVENTION

Renin is a proteolytic enzyme produced and secreted into the bloodstream by the juxtaglomerular cells of the kidney. In the bloodstream, renin cleaves a peptide bond in the serum protein angiotensinogen to produce a decapeptide known as angiotensin I. A second enzyme known as angiotensin converting enzyme, cleaves angiotensin I to produce the octapeptide known as angiotensin II. Angiotensin II is a potent pressor agent responsible for vasoconstriction and elevation of cardiovascular pressure. Attempts have been made to control hypertension by blocking the action of renin or by blocking the formation of angiotensin II in the body with inhibitors of angiotensin I converting enzyme.

Classes of compounds published as inhibitors of the action of renin on angiotensinogen include renin antibodies, pepstatin and its analogs, phospholipids, angiotensinogen analogs, pro-renin related analogs and peptide aldehydes.

5 A peptide isolated from actinomyces has been reported as an inhibitor of aspartyl proteases such as pepsin, cathepsin D and renin [Umezawa et al, in J. Antibiot. (Tokyo), 23, 259-262 (1970)]. This peptide, known as pepstatin, was found to reduce blood pressure in vivo after the injection of hog renin into nephrectomized rats [Gross et al, Science, 175, 656 (1971)]. Pepstatin has the disadvantages of low  
10 solubility and of inhibiting acid proteases in addition to renin. Modified pepstatins have been synthesized in an attempt to increase the specificity for human renin over other physiologically important enzymes. While some degree of specificity has been achieved, this  
15 approach has led to rather high molecular weight hepta- and octapeptides [Boger et al, Nature, 303, 81 (1983)]. High molecular weight peptides are generally considered undesirable as drugs because gastrointestinal absorption is impaired and plasma stability is compromised.


20 Short peptide aldehydes have been reported as renin inhibitors [Kokubu et al, Biochim. Biophys. Res. Commun., 118, 929 (1984); Castro et al, FEBS Lett., 167, 273 (1984)]. Such compounds have a reactive C-terminal aldehyde group and would likely be unstable in vivo.

30 Other peptidyl compounds have been described as renin inhibitors. EP Appl. #128,762, published 18 December 1984, describes dipeptide and tripeptide glyco- containing compounds as renin inhibitors [also see Hanson et al, Biochim. Biophys. Res. Comm., 132, 155-161 (1985), 146, 959-963 (1987)]. EP Appl. #181,110, published 14 May 1986, describes dipeptide histidine derivatives as renin inhibitors. EP Appl. #186,977  
35 published 9 July 1986 describes renin-inhibiting compounds containing an alkynyl moiety, specifically a propargyl glycine moiety, attached to the main chain

between the N-terminus and the C-terminus, such as N-[4(S)-[(N)-[bis(1-naphthylmethyl)acetyl]-DL-propargylglycylamino]-3(S)-hydroxy-6-methylheptanoyl]-L-isoleucinol. EP Appl. #189,203, published 30 July 1986, describes peptidyl-aminodiol as renin inhibitors. EP Appl. #200,406, published 10 December 1986, describes alkyl naphthylmethylpropionyl-histidyl aminohydroxy alkanoates as renin inhibitors. EP Appl. #216,539, published 1 April 1987, describes alkyl naphthylmethylpropionyl aminoacyl aminoalkanoate compounds as renin inhibitors orally administered for treatment of renin-associated hypertension. EP Appl. #229,667, published 22 July 1987, describes acyl aminoacyl aminodiol compounds having a piperazinylcarbonyl or an alkylaminoalkylcarbonyl terminal group at the N-amino acid terminus, such as 2(S)-{[(1-piperazinyl)carbonyl]-oxy}-3-phenylpropionyl}-Phe-His amide of 2(S)-amino-1-cyclohexyl-3(R), 4(S)-dihydroxy-6-methylheptane. PCT Application No. WO 87/04349, published 30 July 1987, describes aminocarbonyl aminoacyl hydroxyether derivatives having an alkylamino-containing terminal substituent and which are described as having renin-inhibiting activity for use in treating hypertension. EP Appl. #300,189 published 25 January 1989 describes amino acid monohydric derivatives having an alkylamino-alkylamino N-terminus and a b-alanine-histidine or sarcosyl-histidine attached to the main chain between the N-terminus and the C-terminus, which derivatives are mentioned as useful in treating hypertension. U.S. Patent No. 4,902,706 which issued 13 February 1990 describes a series of histidineamide-containing amino alkylaminocarbonyl-H-terminal aminodiol derivatives for use as renin inhibitors. U.S. Patent No. 5,032,577 which issued 16 July 1991 describes a series of histidineamide-aminodiol-containing renin inhibitors.


DESCRIPTION OF THE INVENTION

Propargyl glycine amino propargyl diol  
compounds, having utility as renin inhibitors for  
5 treatment of hypertension in a subject, constitute a  
family of compounds of general Formula I:



10 wherein A is selected from methylene, CO, SO and SO<sub>2</sub>;  
wherein X is selected from oxygen atom, methylene and  
15  $\text{>NR}_{10}$  with R<sub>10</sub> selected from hydrido, alkyl and benzyl;  
wherein each of R<sub>1</sub> and R<sub>9</sub> is a group independently  
selected from hydrido, alkyl, cycloalkyl, alkoxyacyl,  
haloalkyl, alkoxy carbonyl, benzyloxy carbonyl,  
loweralkanoyl, haloalkylacyl, phenyl, benzyl, naphthyl,  
and naphthylmethyl, any one of which groups having a  
substitutable position may be optionally substituted  
20 with one or more radicals selected from alkyl, alkoxy,  
alkenyl, alkynyl, halo, haloalkyl, cyano and phenyl, and  
wherein the nitrogen atom to which R<sub>1</sub> and R<sub>9</sub> are  
attached may be combined with oxygen to form an N-oxide;  
wherein R<sub>2</sub> is selected from hydrido, alkyl,  
25 dialkylaminoalkyl, alkylacylaminoalkyl, benzyl and  
cycloalkyl; wherein R<sub>3</sub> is selected from alkyl,  
cycloalkylalkyl, acylaminoalkyl, phenylalkyl,  
naphthylmethyl, aryl, heterocyclicalkyl and  
heterocycliccycloalkyl, wherein the cyclic portion of  
any of said phenylalkyl, naphthylmethyl, aryl,  
30 heterocyclicalkyl and heterocycliccycloalkyl groups may  
be substituted by one or more radicals selected from  
halo, hydroxy, alkoxy and alkyl; wherein each of R<sub>4</sub> and  
R<sub>6</sub> is independently selected from hydrido, alkyl, benzyl


and cycloalkyl; wherein each of R<sub>5</sub> and R<sub>8</sub> is independently selected from



wherein V is selected from hydrido, alkyl, cycloalkyl, haloalkyl, benzyl and phenyl; wherein each of R<sub>13</sub> and R<sub>14</sub> is a radical independently selected from hydrido, alkyl, alkenyl, alkynyl, cycloalkyl, phenyl, heterocyclic, heterocyclicalkyl and heterocycliccycloalkyl; wherein R<sub>7</sub> is selected from substituted or unsubstituted alkyl, cycloalkyl, phenyl, cycloalkylalkyl and phenylalkyl, any one of which may be substituted with one or more groups selected from alkyl, hydroxy, alkoxy, halo, haloalkyl, alkenyl, alkynyl and cyano; wherein each of R<sub>11</sub> and R<sub>12</sub> is independently selected from hydrido, alkyl, haloalkyl, dialkylamino and phenyl; and wherein m is zero or one; wherein n is a number selected from zero through five; wherein p is a number selected from zero through five; and wherein q is a number selected from zero through five; or a pharmaceutically-acceptable salt thereof.

A preferred family of compounds consists of compounds of Formula I wherein A is selected from methylene, CO, SO and SO<sub>2</sub>; wherein X is selected from oxygen atom, methylene and  $\text{>NR}_{10}$  with R<sub>10</sub> selected from hydrido, alkyl and benzyl; wherein each of R<sub>1</sub> and R<sub>9</sub> is independently selected from hydrido, lower alkyl, haloalkyl, cycloalkyl, alkoxycarbonyl, benzyloxycarbonyl, loweralkanoyl, alkoxyacyl, phenyl and benzyl, and wherein the nitrogen atom to which R<sub>1</sub> and R<sub>9</sub>


are attached may be combined with oxygen to form an N-oxide; wherein each of R<sub>2</sub>, R<sub>4</sub> and R<sub>6</sub> is independently selected from hydrido and alkyl; wherein R<sub>3</sub> is selected from phenylalkyl, naphthylmethyl, cyclohexylalkyl, cyclopentylalkyl, heteroarylalkyl and heteroarylcycloalkyl; wherein each of R<sub>5</sub> and R<sub>8</sub> is independently selected from



wherein V is selected from hydrido, alkyl, haloalkyl, benzyl and phenyl; wherein each of R<sub>13</sub> and R<sub>14</sub> is a radical independently selected from hydrido, alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, heteroarylalkyl and heteroarylcycloalkyl; wherein R<sub>7</sub> is selected from substituted or unsubstituted cyclohexylmethyl and benzyl, either one of which may be substituted with one or more groups selected from alkyl, hydroxy, alkoxy, halo and haloalkyl; wherein each of R<sub>11</sub> and R<sub>12</sub> is independently selected from hydrido, alkyl, dialkylamino and phenyl; wherein m is zero or one; wherein n is a number selected from zero through five; wherein p is a number selected from zero through five; and wherein q is a number selected from zero through five; or a pharmaceutically-acceptable salt thereof.

A more preferred family of compounds consists of compounds of Formula I wherein A is selected from methylene, CO, SO and SO<sub>2</sub>; wherein X is selected from oxygen atom, methylene and  $\text{>NR}_{10}$  with R<sub>10</sub> selected from hydrido, alkyl and benzyl; wherein each of R<sub>1</sub> and R<sub>9</sub> is independently selected from hydrido, alkyl, alkoxyacyl,


haloalkyl, alkoxycarbonyl, benzyloxycarbonyl, and  
benzyl, and wherein the nitrogen atom to which R<sub>1</sub> and R<sub>9</sub>  
are attached may be combined with oxygen to form an  
N-oxide; wherein each of R<sub>2</sub>, R<sub>4</sub> and R<sub>6</sub> is independently  
5 selected from hydrido and alkyl; wherein R<sub>3</sub> is selected  
from benzyl, phenethyl, cyclohexylmethyl, phenpropyl,  
pyrrolidinyl, piperidinyl, pyrrolidinylmethyl,  
piperidinylmethyl, pyrazolemethyl, pyrazoleethyl,  
pyridylmethyl, pyridylethyl, thiazolemethyl,  
10 thiazoleethyl, imidazolemethyl, imidazoleethyl,  
thienylmethyl, thienylethyl, furanylethyl,  
furanylethyl, oxazolemethyl, oxazoleethyl,  
isoxazolemethyl, isoxazoleethyl, pyridazinemethyl,  
pyridazineethyl, pyrazinemethyl and pyrazineethyl;  
15 wherein each of R<sub>5</sub> and R<sub>8</sub> is independently selected from



wherein V is selected from hydrido, alkyl and haloalkyl;  
20 wherein each of R<sub>13</sub> and R<sub>14</sub> is a radical independently  
selected from hydrido, alkyl, alkenyl, alkynyl, thiazole  
and thiazolemethyl; wherein R<sub>7</sub> is cyclohexylmethyl;  
wherein each of R<sub>11</sub> and R<sub>12</sub> is independently selected  
from hydrido, alkyl, dialkylamino and phenyl; wherein m  
25 is zero or one; wherein n is a number selected from zero  
through five; wherein p is a number selected from zero  
through five; and wherein q is a number selected from  
zero through five; or a pharmaceutically-acceptable salt  
thereof.


30 An even more preferred family of compounds  
consists of compounds Formula I wherein A is selected  
from CO and SO<sub>2</sub>; wherein X is selected from oxygen atom,

methylene and  $\text{>NR}_{10}$  with  $\text{R}_{10}$  selected from hydrido and  
 methyl; wherein each of  $\text{R}_1$  and  $\text{R}_9$  is independently  
 selected from hydrido, lower alkyl, alkoxyacyl,  
 alkoxy carbonyl, benzyloxycarbonyl, haloalkyl and benzyl,  
 5 and wherein the nitrogen atom to which  $\text{R}_1$  and  $\text{R}_9$  are  
 attached may be combined with oxygen to form an N-oxide;  
 wherein  $\text{R}_2$  is selected from hydrido, methyl, ethyl and  
 isopropyl; wherein  $\text{R}_3$  is selected from benzyl,  
 phenethyl, cyclohexylmethyl, pyrrolidinyl, piperidinyl,  
 10 pyrrolidinylmethyl, piperidinylmethyl, pyrazolemethyl,  
 pyrazoleethyl, pyridylmethyl, pyridylethyl,  
 thiazolemethyl, thiazoleethyl, imidazolemethyl,  
 imidazoleethyl, thienylmethyl, thienylethyl,  
 furanylmethyl, furanylethyl, oxazolemethyl,  
 15 oxazoleethyl, isoxazolemethyl, isoxazoleethyl,  
 pyridazinemethyl, pyridazineethyl, pyrazinemethyl and  
 pyrazineethyl; wherein each of  $\text{R}_4$  and  $\text{R}_6$  is  
 independently selected from hydrido and methyl; wherein  
 each of  $\text{R}_5$  and  $\text{R}_8$  is independently selected from



wherein  $\text{V}$  is selected from hydrido, alkyl and  
 trifluoromethyl; wherein each of  $\text{R}_{13}$  and  $\text{R}_{14}$  is a  
 25 radical independently selected from hydrido, alkyl and  
 alkynyl; wherein  $\text{R}_7$  is cyclohexylmethyl; wherein each of  
 $\text{R}_{11}$  and  $\text{R}_{12}$  is independently selected from hydrido,  
 alkyl, dialkylamino and phenyl; wherein  $m$  is zero;  
 wherein  $n$  is a number selected from zero through five;  
 30 wherein  $p$  is a number selected from zero through five;  
 and wherein  $q$  is a number selected from zero through  
 five; or a pharmaceutically-acceptable salt thereof.

A highly preferred family of compounds consists of compounds of Formula I wherein A is selected from CO and SO<sub>2</sub>; wherein X is selected from oxygen atom and methylene; wherein each of R<sub>1</sub> and R<sub>9</sub> is independently selected from hydrido, methyl, ethyl, n-propyl, isopropyl, benzyl, b, b, b-trifluoroethyl, t-butyloxycarbonyl and methoxymethylcarbonyl, and wherein the nitrogen atom to which R<sub>1</sub> and R<sub>9</sub> are attached may be combined with oxygen to form an N-oxide; wherein R<sub>2</sub> is selected from hydrido, methyl, ethyl and isopropyl; wherein R<sub>3</sub> is selected from benzyl, cyclohexylmethyl, phenethyl, pyrazolemethyl, pyrazoleethyl, pyridylmethyl, pyridylethyl, thiazolemethyl, thiazoleethyl, imidazolemethyl, imidazoleethyl, thienylmethyl, thienylethyl, furanylmethyl, furanylethyl, oxazolemethyl, oxazoleethyl, isoxazolemethyl, isoxazoleethyl, pyridazinemethyl, pyridazineethyl, pyrazinemethyl and pyrazineethyl; wherein each of R<sub>5</sub> and R<sub>8</sub> is independently selected from




wherein V is selected from hydrido, alkyl and trifluoromethyl; wherein each of R<sub>13</sub> and R<sub>14</sub> is a radical independently selected from hydrido, methyl, ethyl, propyl and ethynyl; wherein R<sub>7</sub> is cyclohexylmethyl; wherein each of R<sub>4</sub> and R<sub>6</sub> is independently selected from hydrido and methyl; wherein each of R<sub>11</sub> and R<sub>12</sub> is independently selected from hydrido, alkyl, dialkylamino and phenyl; wherein m is zero; wherein n is a number selected from zero through

five; wherein p is a number selected from zero through five; and wherein q is a number selected from zero through five; or a pharmaceutically-acceptable salt thereof.

5

A more highly preferred class of compounds consists of compounds of Formula I wherein A is selected from CO and SO<sub>2</sub>; wherein X is selected from oxygen atom and methylene; wherein each of R<sub>1</sub> and R<sub>9</sub> is a group independently selected from hydrido, methyl, ethyl, n-propyl, isopropyl, benzyl, b, b, b-trifluoroethyl, t-butyloxycarbonyl and methoxymethylcarbonyl, and wherein the nitrogen atom to which R<sub>1</sub> and R<sub>9</sub> are attached may be combined with oxygen to form an N-oxide; wherein R<sub>2</sub> is selected from hydrido, methyl, ethyl and isopropyl; wherein R<sub>3</sub> is selected from benzyl, cyclohexylmethyl, phenethyl, imidazolemethyl, pyridylmethyl and 2-pyridylethyl; wherein each of R<sub>5</sub> and R<sub>8</sub> is independently selected from

20



wherein V is selected from hydrido, alkyl and trifluoromethyl; wherein each of R<sub>13</sub> and R<sub>14</sub> is a radical independently selected from hydrido, methyl and ethynyl; wherein R<sub>7</sub> is cyclohexylmethyl; wherein each of R<sub>4</sub> and R<sub>6</sub> is independently selected from hydrido and methyl; wherein each of R<sub>11</sub> and R<sub>12</sub> is independently selected from hydrido, alkyl and phenyl; wherein m is zero; wherein n is a number selected from zero through three; wherein p is a number selected from one through three; and wherein q is zero or one; or a pharmaceutically-acceptable salt thereof.

30

T-03034-11201

The term "hydrido" denotes a single hydrogen atom (H). This hydrido group may be attached, for example, to an oxygen atom to form a hydroxyl group; or, as another example, one hydrido group may be attached to a carbon atom to form a  $\text{>CH-}$  group; or, as another example, two hydrido groups may be attached to a carbon atom to form a  $\text{-CH}_2\text{-}$  group. Where the term "alkyl" is used, either alone or within other terms such as "haloalkyl" and "hydroxyalkyl", the term "alkyl"

embraces linear or branched radicals having one to about twenty carbon atoms or, preferably, one to about twelve carbon atoms. More preferred alkyl radicals are "lower alkyl" radicals having one to about ten carbon atoms.

Most preferred are lower alkyl radicals having one to about six carbon atoms. The term "cycloalkyl" embraces cyclic radicals having three to about ten ring carbon atoms, preferably three to about six carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

The term "haloalkyl" embraces radicals wherein any one or more of the alkyl carbon atoms is substituted with one or more halo groups, preferably selected from bromo, chloro and fluoro. Specifically embraced by the term "haloalkyl" are monohaloalkyl, dihaloalkyl and

polyhaloalkyl groups. A monohaloalkyl group, for example, may have either a bromo, a chloro, or a fluoro atom within the group. Dihalalkyl and polyhaloalkyl groups may be substituted with two or more of the same halo groups, or may have a combination of different halo groups. A dihaloalkyl group, for example, may have two fluoro atoms, such as difluoromethyl and difluorobutyl groups, or two chloro atoms, such as a dichloromethyl group, or one fluoro atom and one chloro atom, such as a fluoro-chloromethyl group. Examples of a polyhaloalkyl are trifluoromethyl, 1,1-difluoroethyl, 2,2,2-trifluoroethyl, perfluoroethyl and 2,2,3,3-tetrafluoropropyl groups. The term "difluoroalkyl"

embraces alkyl groups having two fluoro atoms substituted on any one or two of the alkyl group carbon atoms. The terms "alkylol" and "hydroxyalkyl" embrace linear or branched alkyl groups having one to about ten carbon atoms any one of which may be substituted with one or more hydroxyl groups. The term "alkenyl" embraces linear or branched radicals having two to about twenty carbon atoms, preferably three to about ten carbon atoms, and containing at least one carbon-carbon double bond, which carbon-carbon double bond may have either cis or trans geometry within the alkenyl moiety. The term "alkynyl" embraces linear or branched radicals having two to about twenty carbon atoms, preferably two to about ten carbon atoms, and containing at least one carbon-carbon triple bond. The term "cycloalkenyl" embraces cyclic radicals having three to about ten ring carbon atoms including one or more double bonds involving adjacent ring carbons. The terms "alkoxy" and "alkoxyalkyl" embrace linear or branched oxy-containing radicals each having alkyl portions of one to about ten carbon atoms, such as methoxy group. The term "alkoxyalkyl" also embraces alkyl radicals having two or more alkoxy groups attached to the alkyl radical, that is, to form monoalkoxyalkyl and dialkoxyalkyl groups. The "alkoxy" or "alkoxyalkyl" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro or bromo, to provide haloalkoxy or haloalkoxyalkyl groups. The term "alkylthio" embraces radicals containing a linear or branched alkyl group, of one to about ten carbon atoms attached to a divalent sulfur atom, such as a methythio group. Preferred aryl groups are those consisting of one, two, or three benzene rings. The term "aryl" embraces aromatic radicals such as phenyl, naphthyl and biphenyl. The term "aralkyl" embraces aryl-substituted alkyl radicals such as benzyl, diphenylmethyl, triphenylmethyl, phenylethyl, phenylbutyl and diphenylethyl. The terms

"benzyl" and "phenylmethyl" are interchangeable. The terms "aryloxy" and "arylthio" denote radical respectively, aryl groups having an oxygen or sulfur atom through which the radical is attached to a nucleus, examples of which are phenoxy and phenylthio. The terms "sulfinyl" and "sulfonyl", whether used alone or linked to other terms, denotes respectively divalent radicals SO and SO<sub>2</sub>. The term "aralkoxy", alone or within another term, embraces an aryl group attached to an alkoxy group to form, for example, benzyloxy. The term "acyl" whether used alone, or within a term such as acyloxy, denotes a radical provided by the residue after removal of hydroxyl from an organic acid, examples of such radical being acetyl and benzoyl. "Lower alkanoyl" is an example of a more preferred sub-class of acyl. The term "amido" denotes a radical consisting of nitrogen atom attached to a carbonyl group, which radical may be further substituted in the manner described herein. The amido radical can be attached to the nucleus of a compound of the invention through the carbonyl moiety or through the nitrogen atom of the amido radical. The term "alkenylalkyl" denotes a radical having a double-bond unsaturation site between two carbons, and which radical may consist of only two carbons or may be further substituted with alkyl groups which may optionally contain additional double-bond unsaturation.

The term "heterocyclic", as used alone or within groups such as "heterocyclicalkyl" and "heterocycliccycloalkyl", (hereinafter referred to as "heterocyclic-containing groups"), embraces radicals having a saturated, or partially unsaturated, or fully saturated heterocyclic group, wherein the cyclic portion consists of a ring system having one ring or two fused rings, which ring system contains one, two or three hetero atoms as ring members selected from nitrogen, oxygen and sulfur atoms, and which ring system has 4 to

about 12 ring members. Examples of saturated heterocyclic-containing groups are pyrrolidinyl, piperidinyl, pyrrolidinylmethyl, piperidinylmethyl, pyrrolidinylcyclopropyl and piperidinylcyclopropyl. The term "heteroaryl", whether used alone or within the greater terms "heteroarylalkyl" or "heteroarylcycloalkyl", denotes a subset of "heterocyclic-containing groups" having a cyclic portion which is fully-unsaturated, that is, aromatic in character, and which has one or two hetero atoms as ring members, said hetero atoms selected from oxygen, sulfur and nitrogen atoms, and which ring system has five or six ring members. The "heteroaryl" ring may be attached to a linear or branched alkyl radical having one to about ten carbon atoms or may be attached to a cycloalkyl radical having three to about nine carbon atoms. Examples of such heteroarylalkyl or heteroarylcycloalkyl groups are pyrazolemethyl, pyrazoleethyl, pyridylmethyl, pyridylethyl, thiazolemethyl, thiazoleethyl, imidazolemethyl, imidazoleethyl, thienylmethyl, thienylethyl, furanylmethyl, furanylethyl, oxazolemethyl, oxazoleethyl, thiazolylcyclopropyl, imidazolecyclopropyl, thienylcyclopropyl, isoxazolemethyl, isoxazoleethyl, pyridazinemethyl, pyridazineethyl, pyrazinemethyl and pyrazineethyl. The "heterocyclic" portion or "heteroaryl" portion of the radical, as well as the alkyl or cycloalkyl portion of groups containing a "heterocyclic" or "heteroaryl" portion, may be substituted at a substitutable position with one or more groups selected from oxo, alkyl, alkoxy, halo, haloalkyl, cyano, aralkyl, aralkoxy, aryl and aryloxy. Such "heterocyclic", "heterocyclic"-containing group, or "heteroaryl" group may be attached as a substituent through a carbon atom of the hetero ring system, or may be attached through a carbon atom of a moiety substituted on a hetero ring-member carbon

atom, for example, through the methylene substituent of an imidazole-methyl moiety. Also, a heterocyclic or heterocyclic-containing group may be attached through a ring nitrogen atom. For any of the foregoing defined radicals, preferred radicals are those containing from one to about fifteen carbon atoms.

Specific examples of alkyl groups are methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, methylbutyl, dimethylbutyl and neopentyl. Typical alkenyl and alkynyl groups may have one unsaturated bond, such as an allyl group, or may have a plurality of unsaturated bonds, with such plurality of bonds either adjacent, such as allene-type structures, or in conjugation, or separated by several saturated carbons.

0096645-1 RECD  
T0821 545550


Also included in the family of compounds of Formula I are isomeric forms, including diastereoisomers, and the pharmaceutically-acceptable salts thereof. The term "pharmaceutically-acceptable salts" embraces salts commonly used to form alkali metal salts and to form addition salts of free acids or free bases. The nature of the salt is not critical, provided that it is pharmaceutically-acceptable. Suitable pharmaceutically-acceptable acid addition salts of compounds of Formula I may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric and phosphoric acid. Appropriate organic acids may be selected from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic and sulfonic classes of organic acids, example of which are formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, p-hydroxybenzoic, salicylic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, 2-hydroxyethanesulfonic, pantothenic, benzenesulfonic, toluenesulfonic, sulfanilic, mesylic, cyclohexylaminosulfonic, stearic, algenic, b-hydroxybutyric, malonic, galactaric and galacturonic acid. Suitable pharmaceutically-acceptable base addition salts of compounds of Formula I include metallic salts made from aluminium, calcium, lithium, magnesium, potassium, sodium and zinc or organic salts made from N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine. Also included within the phrase "pharmaceutically-acceptable salts" are "quaternary" salts or salts of "onium" cations, such as ammonium, morpholinium and piperazinium cations, as well as any substituted derivatives of these cations where

the salt is formed on the nitrogen atom lone pair of electrons. All of these salts may be prepared by conventional means from the corresponding compound of Formula I by reacting, for example, the appropriate acid or base with the compound of Formula I.

Compounds of Formula I would be useful to treat various circulatory-related disorders. As used herein, the term "circulatory-related" disorder is intended to embrace cardiovascular disorders and disorders of the circulatory system, as well as disorders related to the circulatory system such as ophthalmic disorders including glaucoma. In particular, compounds of Formula I would be useful to inhibit enzymatic conversion of angiotensinogen to angiotensin I. When administered orally, a compound of Formula I would be expected to inhibit plasma renin activity and, consequently, lower blood pressure in a patient such as a mammalian subject (e.g., a human subject). Thus, compounds of Formula I would be therapeutically useful in methods for treating hypertension by administering to a hypertensive subject a therapeutically-effective amount of a compound of Formula I. The phrase "hypertensive subject" means, in this context, a subject suffering from or afflicted with the effects of hypertension or susceptible to a hypertensive condition if not treated to prevent or control such hypertension. Other examples of circulatory-related disorders which could be treated by compounds of the invention include congestive heart failure, renal failure and glaucoma.

**Description of the Synthetic Methods for the  
Preparation of the Renin Inhibitors of the  
Invention**

**Synthetic Scheme 1**




**Synthetic Scheme 1**  
**(Preparation of Compounds of Formula I)**

A suitably protected amino aldehyde 1 is  
5 treated with a Grignard reagent or other organometallic  
reagent, preferably vinylmagnesium bromide, to obtain  
the vinyl carbinol 2. This material, suitably  
protected, is oxidized, preferably with ozone, followed  
by dimethyl sulfide or zinc treatment, to give  
10 intermediate 3. The preceding process is exemplified in  
Hanson, et al., J. Org. Chem. 50, 5399 (1985). This  
aldehyde is reacted with an organometallic reagent such  
as propargylmagnesium bromide to give intermediate 4.  
Other suitable organometallic reagents include  
15 ethynylmagnesium bromide, 1-methylpropynylmagnesium  
bromide, 3-methylpropynyl-magnesium bromide,  
butynylmagnesium bromide and pentynylmagnesium bromide,  
but the choices are not limited to these reagents.  
Compound 4 is deprotected then coupled, using standard  
20 amide/peptide coupling methodology to protected triple  
bond-containing (ethynyl) amino acid derivatives 5 to  
give compound 6. These standard coupling procedures such  
as the carbodiimide, active ester  
(N-hydroxysuccinimide), and mixed carbonic anhydride  
25 methods are shown in Benoiton, et al. J. Org. Chem. 48,  
2939 (1983) and Bodansky, et al. "Peptide Synthesis",  
Wiley (1976). Ethynyl-containing amino acid derivatives  
may be prepared by using procedures such as found in  
Schollkopf, Tetrahedron 39, 2085 (1983). Intermediate 6  
30 is then deprotected, then coupled to intermediate 7  
using the standard amide/peptide coupling methodology,  
to give compounds of Formula I. Suitable protecting  
groups may be selected from among those reviewed by R.  
Geiger in "The Peptides", Academic Press, N.Y. vol. 2  
35 (1979). For example, P<sub>1</sub> may be Boc or Cbz; P<sub>2</sub> may be a  
typical oxygen protective group such as acetyl or t-  
butyldimethylsilyl.

T0221" 54E56650

## Synthetic Scheme 2

Preparation of 7:




## Synthetic Scheme 2

(Preparation of Compounds of Formula I)


Intermediate 7 may be prepared according to the schematic of Synthetic Scheme 2. Intermediate 7 is prepared by coupling amine 8 to mono-protected carboxylic acid 9. Carboxylic acid 9 is a mono-activated moiety by virtue of a suitable leaving group Q which may be chloride, bromide, fluoride, N-hydroxysuccinimido, p-toluenesulfonyloxy or isobutyloxycarbonyloxy, but is not limited to these groups. After coupling, protecting group P<sub>4</sub> is removed (if P<sub>4</sub> is a benzyl group, hydrogenolysis over palladium-on-carbon (Pd-C) is performed) to give intermediate amino acid 7.

### Synthetic Scheme 3


#### Preparation of Intermediate 8



Preparation of **8**, a non-cyclized diamine:



Preparation of **8**:



**Synthetic Scheme 3**  
**(Preparation of Compounds of Formula I)**

Synthetic Scheme 3 describes the preparation of intermediate 8, a non-cyclic diamine. Many of the members of this class, such as ethylene diamine, N,N,N'-trimethylethylene diamine, N,N'-dimethylethylene diamine, N,N'-dimethylpropylene diamine, etc. are commercially available starting materials. Other substituted diamines such as compounds 8a through 8c are obtainable by the procedures depicted in Scheme 3. For example, Boc-L-alanine methyl ester is reduced with diisobutylaluminum hydride to give the corresponding aldehyde which is then reductively aminated with methylamine, then the Boc group is cleaved to give 8a. Alternatively, the procedure of Miller, et al. J. Med. Chem. 19, 1382 (1976) may be employed to give intermediate 8. In another example, a suitably protected diamine is treated with trifluoroacetaldehyde in the presence of sodium cyanoborohydride to give an trifluoroethyl substituent on nitrogen, followed by deprotection to give amine 8.

Abbreviations used:

P1 is an N-protecting group; P2 is H or an oxygen protecting group; P3 is an N-protecting group; P4 is an oxygen protecting group such as benzyl or methyl; Q is a leaving group; Boc is t-butyloxycarbonyl; Cbz is carbobenzoxy.

The following Steps 1-15 constitute specific exemplification of methods to prepare starting materials and intermediates embraced by the foregoing generic synthetic schemes. Those skilled in the art will readily understand that known variations of the conditions and processes of the following preparative procedures can be used to prepare the compounds of Steps 1-15. All temperatures expressed are in degrees Centigrade.

Compounds of Examples 1-99 may be prepared by using the procedures described in the following Steps 1-15:

099645-11201  
FOIA b 7 - D

**Step 1**

(3S,4S)-N-[(tert-butyloxy)carbonyl]-4-amino-3-  
5 (triisopropylsilyloxy)-5-phenylpentene

A solution of (3S,4S)-N-[(tert-butyloxy)carbonyl]-4-amino-3-hydroxy-5-phenylpentene (36.1 mmol), imidazole (90.1 mmol),  
10 chlorotriisopropylsilane (43.3 mmol) and DMF was stirred overnight at room temperature. H<sub>2</sub>O (500ml) was added. The solution was extracted with Et<sub>2</sub>O (2x250 mL). The combined organic material was washed with H<sub>2</sub>O (250 mL), 0.5 M citric acid solution (2x120 mL), dilute KHCO<sub>3</sub>  
15 (2x100 mL), and brine (2x100 mL) and then dried with MgSO<sub>4</sub>. The solvent was evaporated and the residue chromatographed on silica gel to give the title alcohol (12.9 g, 82% yield) <sup>1</sup>H, <sup>13</sup>C, and APT NMR spectral data were consistent with the proposed structure.  
20

**Step 2**

(2R,3S)-N-[(tert-butyloxy)carbonyl]-3-amino-2-  
25 (triisopropylsilyloxy)-4-phenyl-1-butanol

Ozone was bubbled into a solution of the silyl allylic alcohol of Step 1 in CH<sub>2</sub>Cl<sub>2</sub> (244 mL) and MeOH (81 mL) at -78°C until a blue color persisted. The excess ozone was removed with oxygen. To the -78°C  
30 solution, sodium borohydride (80.5 mmol) was added. The mixture was stirred at -78°C for 3 h and then warmed to room temperature and H<sub>2</sub>O (200 mL) was added. The solution was extracted with Et<sub>2</sub>O (3x120 mL). The combined organic layers were washed with brine (2x100  
35 mL) and then dried over Na<sub>2</sub>SO<sub>4</sub>. The filtrate was concentrated to give the title alcohol (13.06g, 100%

yield). The  $^1\text{H}$ ,  $^{13}\text{C}$ , and APT NMR spectral data were consistent with the proposed structure.

Anal. calcd.: C, 63.76; H, 9.43; N, 3.54.  
Found: C, 62.92; H, 9.23; N, 3.47.

5

### Step 3

(2R,3S)-N-[(tert-butyloxy)carbonyl]-3-amino-2-  
(triisopropylsilyloxy)-4-cyclohexyl-1-butanol

10

The monosilylated diol of Step 2 was hydrogenated with 5% Rh-C at 60 psi and 60°C in MeOH (135 mL). The mixture was filtered and the filtrate  
15 evaporated. The residue was purified by chromatography, eluting with (15% EtOAc in hexane) to give the title alcohol (8.70g, 67% yield, mp 61-63°C). The  $^1\text{H}$ ,  $^{13}\text{C}$ , and APT NMR spectral data were consistent with the proposed product.

20

Anal. calcd.: C, 64.96; H, 11.13; N, 3.16.  
Found: C, 65.16; H, 11.26; N, 3.12.

### Step 4

(2R,3S)-N-[(tert-butyloxy)carbonyl]-3-amino-2-  
(triisopropylsilyloxy)-4-cyclohexyl-1-butanal

25

To a -78°C solution of oxalyl chloride (10.8 mmol) in tetrahydrofuran (45 mL) was added  
30 dimethylsulfoxide (21.6 mmol). After stirring for 10 minutes, a solution of the title alcohol of Step 3 (10.8 mmol) in tetrahydrofuran (4.6 mL) was added dropwise. The solution was stirred for 20 minutes at -78°C and then Et<sub>3</sub>N (45.1 mmol) was added. The reaction solution  
35 as allowed to warm to room temperature over a 2 hour period. The opaque, white mixture was poured into water (50 mL) and then extracted with ether (2x50mL). The

combined organic layer was washed with 1N HCl (50 mL), aqueous 5% NaHCO<sub>3</sub> (50 mL), and brine (50 mL). The dried filtrate (MgSO<sub>4</sub>) was evaporated to give the title aldehyde (4.20g, 100% yield). The <sup>1</sup>H, <sup>13</sup>C and APT NMR spectral data were consistent with the proposed structure.

### Step 5

(4RS,5R,6S)-N-[(tert-butyloxy)carbonyl]-6-amino-5-(triisopropylsilyloxy)-7-cyclohexyl-1-heptyn-4-ol

To a solution of magnesium metal (16.4 mmol) and HgCl<sub>2</sub> (0.03g) in Et<sub>2</sub>O (4 mL) in a flask at room temperature was added several drops of 80% propargyl bromide (16.4 mmol) solution in toluene (1.82 mL) contained in an addition funnel. The mixture turned cloudy. Ether (10 mL) was then added to both the flask and to the propargyl bromide solution in the addition funnel. The funnel solution was added dropwise to the reaction mixture which was cooled to 10°C. The white opaque mixture was stirred for 45 minutes at room temperature after the addition was completed. A solution of the title aldehyde of Step 4 (4.64 mmol) in Et<sub>2</sub>O (10 mL) was added dropwise to the flask at room temperature. After 2h at room temperature, the mixture was cooled down to 0°C and a saturated NH<sub>4</sub>Cl solution (25 mL) was added. The Et<sub>2</sub>O layer was collected and the aqueous layer was extracted with Et<sub>2</sub>O (10 mL). The combined organic layers were washed with water (10mL), brine (10 mL) and then dried over MgSO<sub>4</sub>. After evaporation, the residue (a clear, yellow liquid) was purified by medium column chromatography (10% EtOAc in hexane) to give the title alkyne (1.49g, 66.5% yield). The <sup>1</sup>H, <sup>13</sup>C and APT NMR spectral data were consistent for the proposed structure.

## Step 6

(4S,5R,6S)-N-[(tert-butyloxycarbonyl]-6-(amino)-4,5-dihydroxy-7-cyclohexyl-1-heptyne

5 A 1.0M tetra n-butylammonium fluoride solution (21.5 mmol) in THF (21.5mL) was added dropwise to the title alkyne of Step 5 (5.41 mmol) at room temperature. Thin-layer chromatography (TLC) (20% EtOAc in hexane) 10 showed the reaction was completed in 1h. The solution was concentrated and EtOAc (160 mL) was added. The solution was washed with H<sub>2</sub>O (3x60 mL), brine (100 mL), and then dried over MgSO<sub>4</sub>. After evaporation, the residue (a clear, yellow liquid) was purified by column 15 chromatography and the diastereomers were separated. The major isomer (R<sub>f</sub> 0.16, 1.76g, 53% yield) was consistent with the proposed structure from <sup>1</sup>H, <sup>13</sup>C and APT NMR spectral data.

20 Anal. calcd.: C, 66.43; H, 9.60; N, 4.30.  
Found C, 66.00; H, 9.70; N, 4.20.

## Step 7

(4S,5R,6S)-6-amino-7-cyclohexyl-4,5-dihydroxy-1-heptyne

25 To a solution of the diol heptyne of Step 6 isomer (2.85 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (5.5 mL) was added trifluoroacetic acid (71.3 mmol). The reaction was monitored by TLC (50% EtOAc in hexane). After 30 minutes the solution was concentrated and an aqueous 1.0N NaOH 30 solution (7 mL) was added. The solution was extracted with EtOAc (4x10 mL). The organic layer was dried with MgSO<sub>4</sub>. The filtrate was concentrated to give the title amine as a white sticky solid (0.67g, 100% yield). The <sup>1</sup>H, <sup>13</sup>C and APT NMR spectral data were consistent with the 35 prepared structure.

Anal. calc.: C, 68.68; H, 11.08; N, 6.16.  
Found C, 67.58; H, 10.94; N, 5.95.

## Step 8

L-Boc-C-propargylglycine

5

L-C-propargylglycine (10g) [prepared by the method of Schwyzer et al., Helv. Chim. Acta (1976) 59, 2181] was suspended in tetrahydrofuran (30mL). Water (30mL), potassium carbonate (36.7g), and di-*tert*-butyl-dicarbonate (21.9g) were added. Additional water was added to produce a solution which was stirred for 12 hours at room temperature. The organic solvent was then evaporated and the aqueous solution was washed with ether, then acidified to pH 3 with 1N aqueous citric acid. The solution was extracted with methylene chloride and the solvent evaporated to give the title compound (18.9g, 97% yield), used without further purification.

10

15

20

## Step 9

L-Boc-C-propargylglycine amide of (4S,5R,6S)-6-amino-7-cyclohexyl-4,5-dihydroxyhept-1-yne:

25

30

35

L-Boc-C-propargylglycine (1.2g) was dissolved in methylene chloride (5 mL) and N-methyl piperidine (0.57g) was added. The mixture was cooled to zero degrees centigrade and isobutyl chloroformate (0.78g) was added. The mixture was stirred for 10 minutes whereupon the title compound of Step 7 (1.4g) in methylene chloride (5 mL) and *tert*-butyl alcohol (5mL) was added and this mixture stirred for 15 minutes at 0°C and 4°C for 12 hours. The reaction mixture was washed successively with 1N citric acid, saturated sodium hydrogen carbonate, water and brine. The organic layer was dried over magnesium sulfate and evaporated to dryness. The residue was chromatographed on silica gel to give the title compound as a colorless oil. 300 MHz <sup>1</sup>H NMR: consistent with proposed structure.

40

## Step 10

5 L-C-propargylglycine amide of (4S,5R,6S)-6-amino-7-cyclohexyl-4,5-dihydroxyhept-1-yne:

10 The title compound of Step 9 (0.76g) was dissolved in a mixture of trifluoroacetic acid (4.9 mL) and methylene chloride (4.9 mL), and stirred for 30 minutes at room temperature. The solvent was then evaporated and the residue taken up in ethyl acetate. The organic layer was washed with saturated sodium hydrogen carbonate, water and brine, then dried over magnesium sulfate and evaporated to give the title amine.  
15 300 MHz <sup>1</sup>H NMR: consistent with proposed structure.

## Step 11

20 2R-Benzyl butanedioic acid, 1-benzyl ester, dicyclohexylammonium salt

To a slurry of 4-(4-methoxybenzyl) itaconate (prepared by the method of Talley in US Patent # 4,939,288) (50g) in toluene (250mL) was added 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU, 30.4g) in one  
25 portion. Then a solution of benzyl bromide (34.2g) in toluene (50mL) was added dropwise over 0.5 hour. The reaction was stirred for 0.5 hour at room temperature and then poured into a separatory funnel. The mixture was  
30 washed with 3N HCl, aqueous sodium bicarbonate, brine and dried over magnesium sulfate. The solvent was evaporated to give a clear mobile liquid (68g). Chromatography on silica gel, eluting with from 100% hexane to 25% ethyl acetate gave pure 1-(benzyl)-4-(4-methoxybenzyl)  
35 itaconate (55g, 81% yield). A large Fisher-Porter bottle was charged with this itaconate (41g), triethylamine (36g), palladium acetate (380mg), tri-o-tolylphosphine (1.04g) and iodobenzene (24.7g). The bottle was sealed and flushed with nitrogen and placed in an oil bath and

heated for 70 minutes. The residue was chromatographed on silica gel, eluting with 100% hexanes until the less polar impurities were removed. Eluting with 10% ethyl acetate in hexane gave the pure phenyl itaconate. This compound (23.8g) was mixed with toluene (200mL) and the resulting solution treated with trifluoroacetic acid (30mL). The solution was stirred at room temperature for 1.5 hour and then evaporated. The residue was taken up in ether (150mL) and treated with dicyclohexylamine (10.4g) and stirred at 0°C whereupon the salt precipitated. This was isolated by filtration and washed with hexane and dried to give pure 1-benzyl 2-benzylidene succinoate dicyclohexylammonium salt (21.24g, 78% yield). This benzylidene compound (20g) was placed in a Fisher-Porter bottle and also added were degassed methanol (200mL) and rhodium (R,R)DiPAMP (600mg) catalyst. The bottle was sealed and flushed with nitrogen then hydrogen. The reaction was hydrogenated at 40 psig for 15 hours at room temperature. The contents were then poured into a round bottom flask (500mL) and the solvent evaporated to give a dark solid. The residue was taken up in boiling isooctane and allowed to stand, with some title compound crystallizing (7.34g). The non-dissolved residue was taken up in boiling dimethoxyethane. This solution was allowed to cool for 12 hours, whereupon crystals of the title compound formed (6.05g). Combining the two crops gave 13.39g, 66% yield, mp 122-125°C. 300 MHz <sup>1</sup>H NMR: consistent with proposed structure.

30

**Step 12**2R-Benzyl butanedioic acid, 1-benzyl ester

The title compound of Step 11 (9.3g) was suspended in a mixture of water (84mL) and methanol (8.5mL). Solid sodium bisulfate (6.12) was added and the mixture stirred for 5 minutes. The mixture was extracted

with methylene chloride and the combined extracts were dried over magnesium sulfate and evaporated to dryness. The residue was chromatographed on silica gel, eluting with methanol-chloroform-acetic acid (5:95:0.5), to give the pure title compound (4.3g, 74% yield).

### Step 13

Benzyl  $\alpha$ R-[2-[[2-(dimethylamino)ethyl]methylamino]-2-oxoethyl]benzene propanoate

The title compound of Step 12 (4.3g) was dissolved in methylene chloride (20mL) and N-methyl piperidine (1.86g) was added. The mixture was cooled to 0°C and isobutylchloroformate (1.97g) was added. The mixture was stirred for 10 minutes whereupon N,N,N'-trimethylethylene diamine (2.23g) in methylene chloride (10mL) was added. This mixture was stirred at 4°C for 3 hours, then washed with 1N citric acid, saturated aqueous sodium bicarbonate, water and brine. The solvent was evaporated to give the title compound (4.7g, 85% yield). 300 MHz  $^1\text{H}$  NMR: consistent with proposed structure.

### Step 14

$\alpha$ R-[2-[[2-(dimethylamino)ethyl]methylamino]-2-oxoethyl]benzene propanoic acid

The title compound of Step 13 (4.6g) was dissolved in ethanol (50mL) and hydrogenated over 4% Pd-C at 5psi at room temperature for 17 hours. The ethanol was evaporated to give the title compound (3g, 71% yield). 300 MHz  $^1\text{H}$  NMR: consistent with proposed structure.

## Step 15

O-(N-(dimethylaminoethyl)-N-methyl-aminocarbonyl)-3-L-phenyllactic acid

5

10

15

20

Benzyl L-3-phenyllactate (14.28g) was dissolved in tetrahydrofuran (357 mL) and to this was added carbonyl diimidazole (9.78g) and the mixture was stirred at room temperature for 4 hours. N,N,N'-trimethylethylene diamine (6.8g) was added and the mixture stirred for 8 hours. The solvent was evaporated and the residue taken up in ether and washed with water, dried over magnesium sulfate and evaporated to give a yellow oil (13g, 61% yield); 300 MHz <sup>1</sup>H NMR consistent with proposed structure. This ester was hydrogenated over 4% Pd-C @ 5psi and room temperature for 3.5 hours in tetrahydrofuran. The title compound was obtained as a white solid (10g) and recrystallized from methanol.


Anal. calcd for C<sub>15</sub>H<sub>22</sub>N<sub>2</sub>O<sub>4</sub> + H<sub>2</sub>O: C, 57.68; H, 7.75; N, 8.98. Found: C, 57.60; H, 7.82; N, 8.94.

25

30

The following working Examples are provided to illustrate synthesis of Compounds 1-99 of the present invention and are not intended to limit the scope thereof. Those skilled in the art will readily understand that known variations of the conditions and processes of the following preparative procedures can be used to prepare the compounds of the Examples. All temperatures expressed are in degrees Centigrade.


## Example 1



N1-[1R\*-[1S,1R\*-(cyclohexylmethyl)-2S\*,3R\*-dihydroxy-  
hexynyl]amino]carbonyl-3-butynyl-N4-[2-(dimethyl-  
amino)ethyl]-N4-methyl-2S\*-(phenylmethyl)butanediamide

The title compound of Step 14 (0.109g) was dissolved at room temperature in dimethylformamide (2mL) and to this was added N,N'-disuccinimidyl-carbonate (0.96mg), pyridine (148mg) and dimethylaminopyridine (5mg in 0.5mL dimethylformamide). The mixture was stirred for 2 hours, whereupon the title amine of Step 10 (0.12g) was added as a solid in one portion. This mixture was allowed to stir for 12 hours. The solvent was evaporated and the residue dissolved in ethyl acetate. This mixture was washed successively with 5% potassium carbonate, water and brine, then dried over sodium sulfate and the solvent evaporated to afford the title compound (69mg, 31% yield). Proton NMR spectrum: consistent with proposed structure.

## Example 2



[1R\*-[[[[1R\*-[[[[1S,1R\*-(cyclohexylmethyl)-2S\*,3R\*-  
dihydroxy-hexynyl]aminolcarbonyl]-3-  
butynyl]aminolcarbonyl]-2-phenylethyl)]2-  
(dimethylamino)ethyl]methylcarbamate

The title acid of Step 15 is coupled to the  
 title amine of Step 10 using the procedure described for  
 the preparation of Example 1. The crude product is  
 purified by flash chromatography on silica gel, eluting  
 with chloroform-ethanol-ammonium hydroxide (84:15:1), to  
 give title compound.


Compounds #3-99, as shown in Table I below,  
 may be synthesized by reference to the foregoing specific  
 and general procedures for preparing compounds of Formula  
 I.

TABLE I


Example  
Compound No.

Structure


3



4



5




FOIA b 7 - EXEMPT

TABLE I


Example  
Compound No.

Structure


6



7



8




TOBERT SHEPHERD

TABLE I


Example  
Compound No.

Structure


9



10



11







0996345.1.2801

TABLE I

Example  
Compound No.

Structure




TOEAT SHE 9660

TABLE I


Example  
Compound No.

Structure


16



17



18



19



009644-11201  
TOGETHER SHEETS

TABLE I


Example  
Compound No.

Structure


20




21



22



23




T02211 5466550

TABLE I


Example  
Compound No.

Structure


24




25



26



27




TOBERT SHEPHERD

TABLE I


Example  
Compound No.

Structure


28



29



30




Publ. No. 548,966

TABLE I


Example  
Compound No.

Structure


31



32



33




102211 54E9650

TABLE I


Example  
Compound No.

Structure


34



35



36




10221-549550

TABLE I


Example  
Compound No.

Structure

37



38



39







TABLE I

Example  
Compound No.

Structure






FOIEAT SHE9550

TABLE I

Example  
Compound No.

Structure




Downloaded from www.sciencedirect.com

TABLE I


Example  
Compound No.

Structure


46



47



48




T08211" 54E9660

TABLE I


Example  
Compound No.

Structure

49



50



51



0096345-11201  
105211 SHE9660

TABLE I


Example  
Compound No.

Structure


52



53



54




0996345 112001  
FOET SHE5650

TABLE I


Example  
Compound No.

Structure


55



56



57




0996345-11201

TABLE I


Example  
Compound No.

Structure


58



59



60



T03211 54E9660


[illegible]

TABLE I

Example  
Compound No.

Structure

65



66



0999345-112801  
T08211-54E96660

TABLE I


Example  
Compound No.

Structure

67



68



69



090944-1301  
10001-549550

TABLE I


Example  
Compound No.

Structure


70



71



72




T03217 SHE 96560

TABLE I


Example  
Compound No.

Structure

73



74



75








TABLE I

Example  
Compound No.

Structure




09966345.1.2801

TABLE I


Example  
Compound No.

Structure


80




81



82



83




FOI b7E 9660

TABLE I


Example  
Compound No.

Structure


84




85



86



87




TOGETHER SHEETS

TABLE I


Example  
Compound No.

Structure


88



89



90



91


00906445 11201  
TOP SECRET 54E96560




TABLE I


Example  
Compound No.

Structure


96




97



98



99



TOGETHER SHEETS

## BIOLOGICAL EVALUATION

### Human Renin Inhibition in vitro

5           Compounds of Formula I were evaluated as  
inhibitors of human renin in an in vitro assay, as follows:  
This human renin inhibition test has been previously  
described in detail [Papaioannou et al., Clinical and  
Experimental Hypertension, A7(9), 1243-1257 (1985)-]. Human  
10 renin was obtained from the National Institute for  
Biological Standards, London. An incubation mixture was  
prepared containing the following components: in a total  
volume of 0.25mL: 100 mM Tris-acetate buffer at pH 7.4, 25  
x 10<sup>-6</sup> Goldblatt units of renin, 0.05mL of plasma from  
15 human volunteers taking oral contraceptives, 6.0 mM Na-  
EDTA, 2.4 mM phenylmethyl sulfonyl fluoride, 1.5 mM 8-  
hydroxyquinoline, 0.4 mg/mL bovine serum albumin (BSA), and  
0.024 mg/mL neomycin sulfate. This mixture was incubated  
for two hours at 37°C in the presence or absence of renin  
20 inhibitors. The produced angiotensin I was determined by  
radioimmunoassay (New England Nuclear kit). Test compounds  
to be assayed were dissolved in DMSO and diluted with 100mM  
Tris-acetate buffer at pH 7.4 containing 0.5% BSA to the  
appropriate concentration. The final concentration of  
25 organic solvent in the reaction mixture was less than 1%.  
Control incubations at 37°C were used to correct for  
effects of organic solvent on renin activity.

          The in vitro enzymatic conversion of  
30 angiotensinogen to angiotensin I was inhibited by test  
compounds of the invention as indicated in Table II, below:

Table II

Human Renin in vitro  
Inhibition Data

5

Compound Example #                      IC50 Human Renin (nM)

Example 1

34

10

FOOTNOTES: SEE 9550

**Marmoset Plasma Renin Activity (PRA) Inhibition  
on Oral Administration**

The oral activity of renin inhibitor compounds  
5 is determined in vivo in Marmoset monkeys using the  
following procedure. Common Marmoset monkeys (*Callithrix*  
*jacchus*, Charles River, both sexes, body weight 300-400g)  
are placed on a modified high-protein low-sodium diet  
(Purina, St. Louis, MO) for 1 week. About 24 hours prior  
10 to the administration of test compound, Lasix (5mg/kg, im)  
is given. On the day of the test, the animal is  
anesthetized with isoflurane, body weight recorded, and a  
baseline blood sample taken. Then, test compound is given  
intragastrically and blood samples are taken in K-EDTA for  
15 plasma renin activity at appropriate time intervals. The  
PRA is determined by using the protocol outlined below.  
Results may be expressed in terms of PRA at various time  
intervals both before and after compound administrations.  
It is expected that a compound of the invention, when  
20 administered orally, would inhibit marmoset plasma renin  
activity to a level of at about 70% at a dose of 20 mg/kg  
of body weight.

**PLASMA RENIN ACTIVITY ASSAY****I. Angiotensin I Generation**

5	Plasma Sample	200 ul
	PMSF 5%	1 ul
	Neomycin 10%	3 ul
	8-HQ 0.5M	3 ul
	TES 0.5M, pH 7.4	20 ul

10 25 ml of the above reaction mixtures, in duplicate, are incubated at 37°C or 0°C for 2 hours.

**II. Determination of Angiotensin I Concentrations**

Angiotensin I (AI) concentrations are determined by radioimmunoassays with a commercial kit from NEN Co.

**III. Calculation of Plasma Renin Activity**

$$\text{PRA (ng AI/ml/hr)} = (\text{AI at } 37^{\circ} - \text{AI at } 0^{\circ}) / 2 \text{ hr.}$$

25 Abbreviations used:

PMSF : Phenylmethanesulfonylfluoride

8-HQ : 8-Hydroxyquinoline

BSA : Bovine Serum Albumin

TES : N-tris[Hydroxymethyl]methyl-2-aminoethane  
Sulfonic Acid

EDTA : Ethylenediamine Tetraacetic Acid

### Determination of Oral Bioavailability

The bioavailability in marmosets and dogs is determined by sampling the blood via the femoral vein at various time points after administering the renin inhibitor compounds of the invention in polyethylene glycol 400 at an intragastric dose of 10 mg/kg or at an intravenous dose of 1 mg/kg. Compound concentration in plasma is determined using the bioassay technique described below. The percent bioavailability is calculated by dividing the area under the concentration-vs.-time curve obtained from the intragastric experiment by the area under the concentration-vs.-time curve obtained from the intravenous experiment (adjusting for different doses), and multiplying the result by 100%. A compound of the invention would be expected to be orally bioavailable in marmoset and dog.

00996445-12801  
T0821-54E9550

Renin inhibitor plasma concentrations are determined by a bioassay. The plasma samples are extracted with acetonitrile and the extract is evaporated to dryness under nitrogen. Residue is dissolved in 4% bovine serum albumin containing 0.9% NaCl and 5% EDTA. The dissolved residue (0.1 ml) is incubated with a reaction mixture containing human plasma (0.12 ml), 5% phenylmethanesulfonylfluoride (1.2 ul), 10% neomycin (2.4 ul), 0.5 M TES buffer (pH 7.4, 24 ul), and 0.6 mU/ml recombinant human renin (1000 U/mg, 50ul) at 37°C for 90 minutes. The renin activity is determined by a standard angiotensin I radioimmunoassay (New England Nuclear Corp.). The amount of test compound in the plasma is determined by comparing the extent of inhibition of renin activity with that produced by a known amount of test compound added to plasma and analyzed above.

T02211 54E90660

### Renin Inhibitor Species Specificity Method

Blood was collected in a 10 ml Becton Dickinson vacutainer tube with 0.1 ml of 15% EDTA (K3) solution (15 mg) from normal animals or high renin animals; i.e., those pre-treated with 5 mg/kg of Lasix® intramuscularly 2 times within a 6 hour interval 24 hours prior to bleeding. The blood was then centrifuged at 2500 RPM for 20 minutes and the plasma from the various species respectively pooled, aliquoted and stored in the freezer. The human plasma source was a male Caucasian taking a prescription of an angiotensin converting enzyme inhibitor.

The plasma renin activity assay is a modification of the human renin inhibition test in that the animal plasma is the source of both the renin substrate and the renin enzyme. In a total volume of 0.1 ml, 90 mM Tris-acetate buffer, pH 7.5, 12 mM sodium EDTA, 0.012 mg/ml neomycin sulfate, 0.9 mg/ml bovine serum albumin, 1.61 mM phenylmethyl sulfonyl fluoride, 4 mM 8-hydroxyquinoline and 0.09 ml of the animal or human plasmas were incubated for 2 hours at 37°C in a shaking water bath or at 4°C in an ice bath in the presence or absence of renin inhibitors. The produced angiotensin I was determined by radioimmunoassay (New England Nuclear kit).

The amount of angiotensin I generated during the 2 hours at 4°C was usually less than 5% of that activity produced at 37°C; however, the 4°C background values were nevertheless subtracted from the 37°C ones for the 100% activity values. The renin inhibitors are assayed in duplicate using 5 concentrations and the data is expressed as a percent of the 100% total renin activity.

**The Effect of Compound of the Invention on Marmoset  
Blood Pressure Reduction on Intravenous  
Administration**

5           The blood pressure reducing activity of the  
renin inhibitor compound of the invention is determined in  
vivo in Marmoset monkeys using the following procedure.  
Common Marmoset monkeys (*Callithrix jacchus*, Charles River,  
both sexes, body weight ca. 400g) were placed on a  
10 modified high protein low sodium diet (Purina, St. Louis,  
MO) for 1 week. 24 hours prior to the administration of  
test compound, Lasix (5mg/kg, im) was given. On the day of  
the test, the animal was anesthetized with isoflurane, body  
weight recorded, and the left femoral artery and vein were  
15 catheterized. The animal is allowed to regain  
consciousness and the title compound of Example 1 (100  
micrograms/kg) is administered intravenously in 0.1 mL/kg  
polyethylene glycol 400 at time zero. Mean arterial blood  
pressure (MABP) in mmHg is then recorded every ten minutes  
20 until 120 minutes. It is expected that a compound of the  
invention would lower blood pressure in salt-depleted,  
conscious marmosets.

0996345 112804  
FOIA b 7 - DATED 04/25/2011

Also embraced within this invention is a class of pharmaceutical compositions comprising one or more compounds of Formula I in association with one or more non-toxic, pharmaceutically acceptable carriers and/or diluents and/or adjuvants (collectively referred to herein as "carrier" materials) and, if desired, other active ingredients. The compounds of the present invention may be administered by any suitable route, preferably in the form of a pharmaceutical composition adapted to such a route, and in a dose effective for the treatment intended. Therapeutically effective doses of the compounds of the present invention required to prevent or arrest the progress of the medical condition are readily ascertained by one of ordinary skill in the art. The compounds and composition may, for example, be administered intravascularly, intraperitoneally, subcutaneously, intramuscularly or topically.

For oral administration, the pharmaceutical composition may be in the form of, for example, a tablet, capsule, suspension or liquid. The pharmaceutical composition is preferably made in the form of a dosage unit containing a particular amount of the active ingredient. Examples of such dosage units are tablets or capsules. These may with advantage contain an amount of active ingredient from about 1 to 250 mg, preferably from about 25 to 150 mg. A suitable daily dose for a mammal may vary widely depending on the condition of the patient and other factors. However, a dose of from about 0.1 to 3000 mg/kg body weight, particularly from about 1 to 100 mg/kg body weight, may be appropriate.

The active ingredient may also be administered by injection as a composition wherein, for example, saline, dextrose or water may be used as a suitable carrier. A suitable daily dose is from about 0.1 to 100 mg/kg body weight injected per day in multiple doses depending on the

TOBERT SHEPHERD

disease being treated. A preferred daily dose would be from about 1 to 30 mg/kg body weight. Compounds indicated for prophylactic therapy will preferably be administered in a daily dose generally in a range from about 0.1 mg to about 100 mg per kilogram of body weight per day. A more preferred dosage will be a range from about 1 mg to about 100 mg per kilogram of body weight. Most preferred is a dosage in a range from about 1 to about 50 mg per kilogram of body weight per day. A suitable dose can be administered, in multiple sub-doses per day. These sub-doses may be administered in unit dosage forms. Typically, a dose or sub-dose may contain from about 1 mg to about 400 mg of active compound per unit dosage form. A more preferred dosage will contain from about 2 mg to about 200 mg of active compound per unit dosage form. Most preferred is a dosage form containing from about 3 mg to about 100 mg of active compound per unit dose.

The dosage regimen for treating a disease condition with the compounds and/or compositions of this invention is selected in accordance with a variety of factors, including the type, age, weight, sex and medical condition of the patient, the severity of the disease, the route of administration, and the particular compound employed, and thus may vary widely.

For therapeutic purposes, the compounds of this invention are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If administered per os, the compounds may be admixed with lactose, sucrose, starch powder, cellulose esters of alkanolic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin, acacia gum, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets may

contain a controlled-release formulation as may be provided in a dispersion of active compound in hydroxypropylmethyl cellulose. Formulations for parenteral administration may be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions may be prepared from sterile powders or granules having one or more of the carriers or diluents mentioned for use in the formulations for oral administration. The compounds may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or various buffers. Other adjuvants and modes of administration are well and widely known in the pharmaceutical art.

Although this invention has been described with respect to specific embodiments, the details of these embodiments are not to be construed as limitations.